



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# ON THE COMPOSITE.

By PROF. W. H. ECHOLS, Charlottesville, Va.

1. In the first part of my second paper "On Certain Determinant Forms and their Applications" a general form of the Composite (11) was given, which was

$$\begin{vmatrix} fx & , & f_1x & , & \dots , & f_nx & , & f_{n+1}x \\ fy_1 & , & f_1y_1 & , & \dots , & f_ny_1 & , & f_{n+1}y_1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ fy_p & , & f_1y_p & , & \dots , & f_ny_p & , & f_{n+1}y_p \\ Of_{x_1} & , & Of_{1x_1} & , & \dots , & Of_{nx_1} & , & Of_{n+1x_1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ Of_{x_q} & , & Of_{1x_q} & , & \dots , & Of_{nx_q} & , & Of_{n+1x_q} \\ \Phi(u) & , & 0 & , & \dots , & 0 & , & 1 \end{vmatrix} = 0, \quad (a)$$

in which  $O$  was the symbol of differentiation  $D$ , and\*

$$\Phi(u) = \left[ \frac{d}{dx} \right]_{x=u}^{n=p+q} \begin{vmatrix} fx & , & f_1x & , & \dots , & f_nx \\ fy_1 & , & f_1y_1 & , & \dots , & f_ny_1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ fy_p & , & f_1y_p & , & \dots , & f_ny_p \\ Of_{x_1} & , & Of_{1x_1} & , & \dots , & Of_{nx_1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ Of_{x_q} & , & Of_{1x_q} & , & \dots , & Of_{nx_q} \end{vmatrix} \div \left[ \frac{d}{dx} \right]_{x=u}^{n=p+q} \begin{vmatrix} f_1x & , & \dots , & f_{n+1}x \\ f_1y_1 & , & \dots , & f_{n+1}y_1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ f_1y_p & , & \dots , & f_{n+1}y_p \\ Of_{1x_1} & , & \dots , & Of_{n+1x_1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ Of_{1x_q} & , & \dots , & Of_{n+1x_q} \end{vmatrix}. \quad (b)$$

2. I propose now to show that the same formula holds good when  $O$  is the symbol  $J$  of Finite Differences.

In my first paper, at the bottom of page 109, I deduced the formula

$$h^n f^n(u) = fx - C_{n,1} f(x-h) + \dots + c_{n,r} f(x-rh) + \dots + (-1)^n f(x-nh),$$

$C_{n,r}$  representing the binomial coefficient, and  $u$  some value of  $x$  lying between  $x$  and  $x-nh$ .

---

\* In (11),  $\left[ \frac{d}{dx} \right]_{x=u}^{n=p+q}$  was given as  $\left[ \frac{d}{dx} \right]_{x=u}^{q+1}$ . It is so easy to extend the operation to  $n = p+q$  that we do not give it here.

The second member of this equality is the well known interpolation formula of Finite Differences (Boole's Finite Differences, p. 19), and its value is known to be  $\Delta^n f x$ . We have, therefore,

$$\Delta^n f x = h^n f^n(u). \quad (c)$$

Lagrange's form of Rolle's theorem, written in this notation is

$$\Delta f x = h f'(u).$$

Therefore the above may be regarded as the proper generalization of Lagrange's form, a formula which we now require in our analysis.

Let  $O \equiv \Delta$ , the symbol of Finite Differences. Let  $M$  and  $N$  be the minors of 1 and  $\Phi(u)$  in (a), respectively. Put

$$M = (-1)^n N R,$$

where  $R$  is some unknown function of  $x$ , and let this equation become

$$M_0 = (-1)^n N_0 R_0,$$

wherein we have substituted for the variable  $x$  some arbitrary constant  $x_0$ . Consider the function

$$F = M - (-1)^n N R_0.$$

We have  $F = 0$  when  $x = x_0, y_1, \dots, y_p$ ; therefore the first derivative  $F' = 0$  for  $p$  values of  $x$ , say  $u_1, \dots, u_p$ , lying respectively between each pair of values  $x_0, y_1$ ;  $y_1, y_2$ ;  $\dots$ ;  $y_{p-1}, y_p$ .

Now  $\Delta F = 0$  when  $x = x_1$ . Hence (if the scale of difference be  $h$ ), since we have  $\Delta F x = h F'(u)$  ( $u$  lying between  $x$  and  $x + h$ ), we must have  $F' = 0$  for some value of  $x$  between  $x_1$  and  $x_1 + h$ , say  $x_1 + h_1$ . Therefore  $F'$  vanishes  $p + 1$  times at the values indicated. It follows, therefore, that  $F''$  vanishes  $p$  times, once between each pair of values  $u_1, u_2$ ;  $\dots$ ;  $u_{p-1}, u_p$ ;  $u_p$  and  $x_1 + h_1$ , say when  $x = u'_1, \dots, u'_p$ .

Again, since we have

$$\Delta^2 F x = h^2 F''(u)$$

( $u$  between  $x$  and  $x + 2h$ ), and since  $\Delta^2 F = 0$  when  $x = x_2$ , then must  $F'' = 0$  for some value of  $x$  between  $x_2$  and  $x_2 + 2h$ , say  $x_2 + h_2$ .  $F''$  therefore vanishes  $p + 1$  times for values of  $x$  lying between the pairs of values  $u'_1, u'_2$ ;  $\dots$ ;  $u'_{p-1}, u'_p$ ;  $u'_p$  and  $x_2 + h_2$ .

Reasoning in the same way, we proceed until we show that the  $q$ th derivatives of  $F$  vanishes  $p + 1$  times for values of  $x$  which lie between determinate limits; and that, finally, the  $(p + q)$ th or  $n$ th derivative of  $F$  vanishes once for some value of  $x$ , say  $u$ , which lies between the greatest and least of the

quantities

$$x_0, y_1, \dots, y_p, x_1 + h, \dots, x_q + qh;$$

so that we have

$$F_{x=u}^{n=p+q} = M_{x=u}^{n=p+q} - (-1)^n N_{x=u}^{n=p+q} R_0 = 0.$$

If we put

$$\Phi(u) = M_{x=u}^{n=p+q} / N_{x=u}^{n=p+q},$$

then

$$M_0 = (-1)^n \Phi(u) N_0;$$

and since  $x_0$  is arbitrary we may drop the subscript and write

$$M = (-1)^n \Phi(u) N,$$

which is formula (a).

It is to be observed that when in (a),  $O \equiv D$ , the symbol of differentiation,  $u$  lies between the greatest and least of  $x, x_1, \dots, x_q, y_1, \dots, y_p$ .

3. The rationale of the process illustrating the application of the composite for differentiation to the expression of functions in an infinite series of other functions may be presented thus :—

Let there be two functions  $f(x)$  and

$$\sum_0^{n-1} A_r \varphi_r x = A_0 + A_1 \varphi_1 x + \dots + A_{n-1} \varphi_{n-1} x.$$

Let the difference between these two functions be  $R$ , so that

$$fx = A_0 + A_1\varphi_1x + \dots + A_{n-1}\varphi_{n-1}x + R. \quad (c)$$

Let  $a_1, \dots, a_n$  be certain arbitrary values of the variable  $x$ , and let us have

$$\left. \begin{aligned} fa_1 &= A_0 + A_1\varphi_1a_1 + \dots + A_{n-1}\varphi_{n-1}a_1 + R_1, \\ . & . . . . . \\ fa_n &= A_0 + A_1\varphi_1a_n + \dots + A_{n-1}\varphi_{n-1}a_n + R_n. \end{aligned} \right\} \quad (\text{d})$$

In these  $n$  relations (d) there are  $n$  undetermined arbitrary quantities  $A_0, A_1, \dots, A_{n-1}$ . Let us determine these so that we shall have  $R_1 = 0, \dots, R_n = 0$ . Thus, the value of  $A_r$  which satisfies this condition is

$$A = (-1)^r \frac{\begin{vmatrix} f_{a_1}, 1, \varphi_1 a_1, \dots, \varphi_{r-1} a_1, \varphi_{r+1} a_1, \dots, \varphi_{n-1} a_1 \\ \vdots \\ f_{a_n}, 1, \varphi_1 a_n, \dots, \varphi_{r-1} a_n, \varphi_{r+1} a_n, \dots, \varphi_{n-1} a_n \end{vmatrix}}{\begin{vmatrix} 1, \varphi_1 a_1, \dots, \varphi_{n-1} a_1 \\ \vdots \\ 1, \varphi_1 a_n, \dots, \varphi_{n-1} a_n \end{vmatrix}}.$$







$$\left. \begin{aligned} f(y+z_0) &= A_0 + A_1\varphi_1(y+z_0) + \dots + A_{n-1}\varphi_{n-1}(y+z_0) + R_0 \\ f(y+2z_1) &= A_0 + A_1\varphi_1(y+2z_1) + \dots + A_{n-1}\varphi_{n-1}(y+2z_1) + R_1 \\ &\vdots \\ f(y+nz_{n-1}) &= A_0 + A_1\varphi_1(y+nz_{n-1}) + \dots + A_{n-1}\varphi_{n-1}(y+nz_{n-1}) + R_{n-1} \end{aligned} \right\} \quad (\text{II})$$

In the  $n$  relations II we have  $n$  undetermined quantities  $A_0, \dots, A_{n-1}$ . Let us determine these so that the functions  $f_x$  and  $\Sigma A_r \varphi_{r,x}$  shall coincide for the  $n$  values  $x = y + z_0, \dots, y + nz_{n-1}$ , or, what is the same thing, so that  $R_0 = 0, \dots, R_{n-1} = 0$ .

The value of  $A_r$ , which satisfies this condition is

$$1) \frac{\begin{vmatrix} f(y+z_0) & 1, \varphi_1(y+z_0) & \dots, \varphi_{r-1}(y+z_0) & \varphi_{r+1}(y+z_0) & \dots, \varphi_{n-1}(y+z_0) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ f(y+nz_{n-1}) & 1, \varphi_1(y+nz_{n-1}), \dots, \varphi_{r-1}(y+nz_{n-1}), \varphi_{r+1}(y+nz_{n-1}), \dots, \varphi_{n-1}(y+nz_{n-1}) \end{vmatrix}}{\begin{vmatrix} 1, \varphi_1(y+z_0) & \dots, \varphi_{n-1}(y+z_0) \\ \vdots & \vdots \\ 1, \varphi_1(y+nz_{n-1}), \dots, \varphi_{n-1}(y+nz_{n-1}) \end{vmatrix}}. \quad (\text{III})$$

Let us consider the  $A$ 's in I and II to have these values, then

$$\frac{\left| \begin{array}{cccc} f^x & , 1, \varphi_1 x & , \dots, \varphi_{n-1} x \\ f(y+z_0) & , 1, \varphi_1(y+z_0) & , \dots, \varphi_{n-1}(y+z_0) \\ \cdot & \cdot & \cdot & \cdot \\ f(y+nz_{n-1}) & , 1, \varphi_1(y+nz_{n-1}), \dots, \varphi_{n-1}(y+nz_{n-1}) \end{array} \right|}{\left| \begin{array}{cccc} 1, \varphi_1(y+z_0) & , \dots, \varphi_{n-1}(y+z_0) \\ \cdot & \cdot & \cdot & \cdot \\ 1, \varphi_1(y+nz_{n-1}), \dots, \varphi_{n-1}(y+nz_{n-1}) \end{array} \right|} = R. \quad (\text{IV})$$

Let  $nz = y - w$ ,  $y$  and  $w$  being fixed definite values of  $x$ . Let  $z_0, z_1, \dots, z_{n-1}$  approach the value  $z$ , and therefore the value zero when  $n$  approaches infinity. Then when  $n = \infty$ , the functions  $f_x$  and  $\sum A_{r,\varphi_r}x$  coincide in an infinite number of consecutive values of  $x$  between  $y$  and  $w$ . The above value of  $R$ , however, in the limit takes the indeterminate form  $0/0$ . For any finite number of rows after the first become identical, the elements in the  $r$ th column in these rows being all  $\varphi_r y$ . Also, any finite number of rows from the bottom are identical, the elements in the  $r$ th column in these rows all being  $\varphi_r w$ . To



remove this indeterminateness, we apply the operator

$$\left[ \frac{d}{dz_1} \right]_{z_1=z}^1 \cdots \left[ \frac{d}{dz_{n-1}} \right]_{z_{n-1}=z}^{n-1},$$

to the numerator and denominator and divide by  $2^1 \cdot 3^2 \cdots n^{n-1}$ . We can now make  $z = 0$  by making  $n = \infty$  without indetermination.

Thus, we have

$$\begin{vmatrix} fx & , & 1, \varphi_1 x & , & \dots, \varphi_{n-1} x \\ f(y+z) & , & 1, \varphi_1(y+z) & , & \dots, \varphi_{n-1}(y+z) \\ f'(y+2z) & , & 0, \varphi_1'(y+2z) & , & \dots, \varphi_{n-1}'(y+2z) \\ \dots & & \dots & & \dots \\ f^{r-1}(y+rz) & , & 0, \varphi_1^{r-1}(y+rz) & , & \dots, \varphi_{n-1}^{r-1}(y+rz) \\ \dots & & \dots & & \dots \\ f^{n-r-1}(y+n-rz), 0, \varphi_1^{n-r-1}(y+n-rz), \dots, \varphi_{n-1}^{n-r-1}(y+n-rz) \\ \dots & & \dots & & \dots \\ f^{n-1}(y+nz) & , & 0, \varphi_1^{n-1}(y+nz) & , & \dots, \varphi_{n-1}^{n-1}(y+nz) \end{vmatrix} = R. \quad (V)$$

$$\begin{vmatrix} \varphi_1'(y+z) & , & \dots, \varphi_{n-1}'(y+z) \\ \dots & & \dots \\ \varphi_1^{n-1}(y+nz), \dots, \varphi_{n-1}^{n-1}(y+nz) \end{vmatrix}$$

If now the coefficient of  $\varphi_r x$  in the expansion of this determinant be determinate when  $n = \infty$ , so that the series  $\sum A_{\varphi_r x}$  is convergent; then, when  $n = \infty$ , and  $R$  is zero when  $x$  lies between  $y$  and  $w$ , the functions  $fx$  and  $\sum A_{\varphi_r x}$  coincide for all values of  $x$  between  $y$  and  $w$ . The composite shows the value of  $R$  to be also

$$\begin{vmatrix} 1, \varphi_1 x & , & \dots, \varphi_n x \\ 1, \varphi_1(y+z) & , & \dots, \varphi_n(y+z) \\ \dots & & \dots \\ 0, \varphi_1^{n-1}(y+nz), \dots, \varphi_n^{n-1}(y+nz) \end{vmatrix} \cdot \begin{vmatrix} f^n u & , & \varphi_1^n u & , & \dots, \varphi_n^n u \\ f''(y+2z) & , & \varphi_1''(y+2z) & , & \dots, \varphi_{n-1}''(y+2z) \\ \dots & & \dots \\ f^{n-1}(y+nz), \varphi_1^{n-1}(y+nz), \dots, \varphi_{n-1}^{n-1}(y+nz) \end{vmatrix}, \quad (V)$$

$$\begin{vmatrix} \varphi_1'(y+z) & , & \dots, \varphi_{n-1}'(y+z) \\ \dots & & \dots \\ \varphi_1^{n-1}(y+nz), \dots, \varphi_{n-1}^{n-1}(y+nz) \end{vmatrix} \cdot \begin{vmatrix} \varphi^n u & , & \varphi_2^n u & , & \dots, \varphi_n^n u \\ \varphi_1'(y+2z) & , & \varphi_2'(y+2z) & , & \dots, \varphi_n'(y+2z) \\ \dots & & \dots \\ \varphi_1^{n-1}(y+nz), \varphi_2^{n-1}(y+nz), \dots, \varphi_n^{n-1}(y+nz) \end{vmatrix}$$

